



# UNITED STATES PATENT AND TRADEMARK OFFICE

UNITED STATES DEPARTMENT OF COMMERCE  
United States Patent and Trademark Office  
Address: COMMISSIONER FOR PATENTS  
P.O. Box 1450  
Alexandria, Virginia 22313-1450  
www.uspto.gov

APPLICATION NO.	FILING DATE	FIRST NAMED INVENTOR	ATTORNEY DOCKET NO.	CONFIRMATION NO.
10/595,821	01/16/2007	August Verbruggen	50304/021002	7212
21559	7590			
CLARK & ELBING LLP				
101 FEDERAL STREET				
BOSTON, MA 02110				
EXAMINER				
ARIANE, KADE				
ART UNIT		PAPER NUMBER		
1651				
NOTIFICATION DATE		DELIVERY MODE		
05/27/2010		ELECTRONIC		

**Please find below and/or attached an Office communication concerning this application or proceeding.**

The time period for reply, if any, is set in the attached communication.

Notice of the Office communication was sent electronically on above-indicated "Notification Date" to the following e-mail address(es):

patentadministrator@clarkelbing.com

Attachment to the Advisory Action:

1

Applicant argues that because Verbruggen teaches xylosan polysulfate and chondroitin polysulfate but not heparin, significantly increase aggrecan aggregate sizes, therefore Verbruggen teaches away from using heparin-like molecules interchangeably in chondrocyte cultivation. Applicant argues that Verbruggen concludes that heparin-like molecules have different or unpredictable effects on chondrocyte development (see Remarks page 3 2<sup>nd</sup> -4<sup>th</sup> paragraphs filed on 05/04/2010). These arguments are considered but are not found persuasive because Verbruggen et al. do not teach xylosan polysulfate and chondroitin polysulfate are heparin-like molecules (or heparinoids), instead Verbruggen et al. teach xylosan polysulfate, chondroitin polysulfate are polysulphated polysaccharides. Verbruggen et al. teach polysulphated polysaccharides, xylosan polysulfate, chondroitin polysulfate, increased the synthesis of high molecular weight hyaluronan by chondrocyte. Verbruggen et al. further teach stimulation of the production of high molecular weight hyaluronan by chondrocytes seems to be a common effect of polysulfated polysaccharides (p1669 1<sup>st</sup> column 5<sup>th</sup> paragraph lines 24-28).

Moreover, Verbruggen et al. teach polysulphated polysaccharides, xylosan polysulfate and chondroitin polysulfate, consistently improved aggrecan synthesis in the culture chondrocytes, and heparin was less active on these cells (page 1669 1<sup>st</sup> column 4<sup>th</sup> paragraph lines 1-4). Therefore, Verbruggen et al. neither teach away from using heparin (or heparin-like molecules) in chondrocyte cultivation nor concludes that heparin-like molecules have unpredictable effects on chondrocyte development, as alleged by the Applicant.

Applicant argues that one of ordinary skill in the art would have not been motivated to combine Verbruggen and Rosenberg teachings and that the combination of references fails to provide a motivation to use polysulphated alginate for chondrocyte cultivation (see Remarks page 3 4<sup>th</sup> paragraphs filed on 05/04/2010). These arguments are considered but are not found persuasive because Rosenberg et al. teach polysulfates prepared from alginic acid (polysulphated alginate) and chondroitin polysulphates are polysulfated polysaccharides, which are non-thrombogenic (heparin-like effect). Rihova teach a matrix suitable for implantation, must be non-thrombogenic (anticoagulant activity) to be biocompatible. Therefore, it would have been obvious to a person of ordinary skill in the art at the time the invention was made to try (choosing from a finite number of identified polysulfated polysaccharides) and to use polysulfated polysaccharide, polysulphated alginate, in the method and the composition as taught by Verbruggen et al. with a reasonable expectation of success in providing an in vitro method for cultivation of chondrogenic cells, a composition comprising polysulphated alginate, and a method of treatment of cartilage defects. The motivation as taught by Verbruggen et al. would be stimulation of the production of high molecular weight hyaluronan by chondrocytes seems to be a common effect of polysulfated polysaccharides.